

Chemical Composition and Insecticidal Activity of *Laurus nobilis* Essential Oil on *Culiseta longiareolata* (Diptera : Culicidae) larvae

Oulfa BOUZIDI, Fouzia TINE-DJEBBAR, Samir TINE and Noureddine SOLTANI

Abstract—The current study was undertaken in order to determine larvicidal activity of *Laurus nobilis* essential oil and its effects at LC25 and LC50 on biochemical composition and on the body weight of the fourth instar larvae of *Culiseta longiareolata* (Diptera: Culicidae), the most common and abundant species in Tebessa area (Northeast Algeria). The obtained percent yield of the hydrodistilled oil from aerial parts of *Laurus nobilis* was $0.96 \pm 0.045\%$. The GC/MS analysis of *Laurus nobilis* essential oil has led to the identification of 56 components. Eucalyptol (25.62 %), Linalool (11.83 %), Methyl Eugenol (11.07%) and Camphene (10.18%), were the major constituents of which. Bioassay test done following the World Health Organization standard protocol revealed that this essential oil exhibited larvicidal activity with dose-response relationship. The morphometric study shows that the essential oil tested was found to decrease the growth of larvae. Moreover, it reduces significantly the body contents of proteins, carbohydrates and lipids of treated individuals. Overall, our results indicate that *L. nobilis* essential oil has potential for the development of new and safe control products against mosquitoes.

Keywords— *Culiseta longiareolata*, Essential oil, *Laurus nobilis*, Toxicity, Chemical composition

I. INTRODUCTION

Insects are a very important part of the biodiversity terrestrial and aquatic ecosystems. Hematophagous insects like mosquitoes play an important role in global disease transmission such as Zika, dengue fever, yellow fever, chikungunya, malaria and Japanese encephalitis [1]. Mosquito controls, using synthetic chemical insecticides have adverse effects on the environment and also cause growing of insecticide resistance in arthropods [2-3]. However, the search for new insect control agents from natural products which are selective, biodegradable and of low environmental toxicity is crucial [4]. Mosquito larvae can be controlled by natural enemies [5] such as larvivorous fish [6], bacteria [7] and by insect growth disruptors (IGDs) [8-9]. The phytochemicals derived from plant resources can act as larvicides, insect growth regulators, repellents, and ovipositional attractants, having deterrent activities observed by different researchers [10-12].

In general, essential oils have been considered as important

Oulfa BOUZIDI, Ph.D. Student, University of Larbi Tebessi, TEBESSA, Algeria.

Fouzia TINE-DJEBBAR, Samir TINE and Noureddine SOLTANI, University of Larbi Tebessi, TEBESSA, Algeria.

natural resources to act as insecticides [13-14], with low mammalian toxicity and rapidly degradable in the environment [15]. They show a various bioactivities against mosquito species with ovicidal, larvicidal, pupicidal [16-17] and adulticidal potentials [18-19].

Approximately, 2000 plant species have been known to produce secondary metabolites of value in biological pest control programs and only 344 plant species showed insecticidal activity against mosquitoes [20-23]. Essential oils can be synthesized by all plant organs (flowers, buds, seeds, leaves, twigs, bark, herbs, wood, fruits and root) and can therefore be extracted from these parts [24]. They are mainly formed by mixtures of monoterpenes, sesquiterpenes, phenylpropanoids and metabolites that confer the mixtures with organoleptic characteristics and biological activities [25]. *Laurus nobilis* L. is evergreen shrub, belonging to family Lauraceae comprises 32 genera and about 2000-2500 species. Some studies have shown that the Laurel essential oil used as antioxidant [26], antifungal [27], antibacterial [28] and insecticidal agent [29]. *Culiseta longiareolata* M. represent the most representative mosquito species in the Tebessa area [30]. The present study was designed to determine the chemical composition of the *L. nobilis* essential oil and its larvicidal activity against fourth instar larvae of *C. longiareolata*. In addition, its effects on morphometric measurements and on main biochemical components (carbohydrates, proteins and lipids) in whole body were investigated.

II. DETAILS EXPERIMENTAL

A. Plant materials and extraction of the essential oil

The fresh aerial parts of *L. nobilis* samples were collected in Tebessa area (Northeast Algeria) in march-july 2016. Dried leaves of the plants (about 100 g) were cut into small pieces and hydrodistilled in a clevenger type apparatus for 3 h according to the method recommended in the British Pharmacopoeia (1988). The volatile oils were dried over anhydrous sodium sulfate and stored in sealed vials at 4 °C until analysis. The yield of the oils was calculated based on dried weight of plant.

B. Gas chromatography-mass spectrometry

The GC/MS analysis were performed with an HP Agilent 2890 plus gas chromatograph (GC) equipped with a HP-5MS column (a length of 30 m × internal diameter of 0.25 mm, and 0.25 mm film thickness). The helium was used as transporter gas. The GC oven temperature was kept at 60°C for 8 min and programmed to 250°C for 10 min at rate of 2°C/min. The

injector temperature was set at 250 °C. The split flow was adjusted at 50ml/min. MS were taken at 70 eV. The sample was dissolved in pure hexane. A volume of 0.2µl was injected for GC-MS analysis. Constituent's identification was found on comparison of retention times with those of corresponding reference standards using the NIST 02 and WILEY 7N libraries [31; 32]. Percentage compositions of essential oil were calculated according to the area of the chromatographic peaks.

C. Mosquito rearing

The larvae of *C. longiareolata* (Diptera: Culicidae) were obtained from a stock colony of the laboratory. Each 20 larvae were kept in Pyrex storage jar containing 150 ml of stored tap water and they were daily fed with fresh food consisting of a mixture of Biscuit Petit Regal-dried yeast (75:25 by weight). The water was replaced every three days.

D. Larvicidal test

Newly ecdysed fourth-instar larvae of *C. longiareolata* were exposed to the different concentrations (25, 50, 100, and 200 ppm) for 24 h. According to the World Health Organization, the larvae were removed, and placed in clean water. The test was realized with three repetitions containing 20 larvae each. Mortality was registered at 1st day, 3rd day, 5th day and 7th day following treatment. The control mortality was corrected by Abbott's formula. Sub-lethal and lethal concentrations (LC₂₅, LC₅₀ and LC₉₀) and 95% confidence limits (95% CL) were calculated.

E. Morphometric measurements

As above, newly molted fourth instar larvae were treated with *L. nobilis* EO at its LC₂₅ and LC₅₀ as determined before. The body size was recorded by measuring the width of the thorax in larvae [33] on 3 replicates of 20 individuals and the body weight was also determined.

F. Biochemical procedure

Carbohydrates, Proteins and lipids were extracted following the procedure of Shibko et al. (1966) [34]. Pooled samples (20 individuals per pool) were weighed and extracted in 1 ml of trichloroacetic acid (20%). In brief, quantification of proteins was carried following the Coomassie Brilliant Blue G-250 dye-binding method [35] with bovine serum albumin as a standard. The absorbance was measured at 595 nm. Carbohydrates were determined following the anthrone method [36] using glucose as standard. Lipids were measured by the vanillin method [37]. Data were expressed in µg per individual.

G. Statistical analysis

Results are presented as mean ± standard deviation (SD). The significance between different series was tested using Student's t tests at 5 % level. Data were subjected to one-way analysis of variance (ANOVA) followed by a post-hoc honestly significant difference (HSD) Tukey's test.

III. RESULTS AND DISCUSSION

A. Extraction yield and chemical composition of essential oil

The results of the steam distillation show that the yield of *L. nobilis* EO was 0.45± 0.96% (dry leaves of the plant). The percentage and the retention time of the identified compound of

this essential oil were given in Table 1 and figure 1. Gas chromatography and mass spectrometry analysis resulted in the identification of fifty-six compounds, with Eucalyptol (25.62 %) and Linalool (11.83 %) as the major components. The variations of the chemical components of *L. nobilis* essential oil related to the geographic origins, genetic variability, growing conditions, organ development, seasonal, methods of extraction and the plant part from which it was extracted such as seed, leaf, flower and fruits [38]. Compared to the published data, the chemical profile obtained presented differences, but also some similarities.

This difference is confirmed by the chemical composition of essential oil of fresh bay leaves from Turkey with 1,8 cineol (64.61%) as a major component. It has been found that the *L. nobilis* oil from Algeria contains major components, 1-8 cineol (39.69%) and camphene (14.21%) [39]; from Brazil; 1-8 cineol ((35.50 g/100 g) and linalool (14.10 g/100 g) [40], from Turkey; 1,8-cineole (51.73-68.48%) and α-terpinyl acetate (4.04-9.87%) [41] and from Tunisia ; 1,8-cineole (56%) and α-terpinyl acetate (9.0%). Mediouni et al. (2012)[42] mentioned that three *L. nobilis* EO from Algeria, Tunisia and Morocco showed quantitative, rather than qualitative, differences in their chemical composition that depended on their cultivation locations.

TABLE I: CHEMICAL COMPOSITION OF *L. NOBILIS* OIL: RETENTION TIME (RT) AND AREA (%) OF 10 MAJOR CONSTITUENTS

N°	RT	Compounds	Area %
1	14.14	Eucalyptol	25.62
2	19.35	Linalool	11.83
3	39.92	Methyl Eugenol	11.07
4	36.00	α-terpinyl acetate	10.18
5	10.28	Sabinene	7.34
6	7.98	α-Pinene	3.87
7	25.35	1-α-Terpineol	3.45
8	40.03	Veratrole methyl	2.73
9	36.62	Eugenol	1.88
10	55.93	Shyobunol	1.45




Fig. 1. GC-MS chromatogram for essential oil obtained from *L. nobilis* (abundance as function the time in min).

B. Insecticidal activity

Essential-oil extracted from plants deed as promising alternative natural Products for the control of many insect pests particularly mosquitoes [43]. Dose-response relationship was

determined for *L. nobilis* essential oil applied to newly molted fourth instar larvae of *C. longiareolata* (Table 2 and Fig. 2). Preliminary tests showed that the solvent (ethanol) had no significant effect on larvae compared to untreated series. The mortality was scored at different periods after treatment (1st, 3rd, 5th and 7th day). After the treatment, the intoxicated larvae showed a change in their behaviour by sinking to the bottom of the jar and remain there motionless until they died. Our results indicate that the *L. nobilis* essential oil and its active components could be developed as control agents against the mosquito larvae. Diverse studies reported the larvicidal activity of monoterpenes against various species of mosquitoes [44-45].




Fig. 2. Efficacy of *L. nobilis* EO against fourth instar larvae of *C. longiareolata*

TABLE II. LARVICIDAL ACTIVITY OF *L. NOBILIS* EO AGAINST *C. LONGIAREOLATA* AT DIFFERENT PERIODS AFTER TREATMENT.

Time (Day)	Slope	R ²	CL ₂₅ (ppm)	CL ₅₀ (ppm)	CL ₉₀ (ppm)
			95% FL	95% FL	95% FL
1	0.34	0.98	132.80	203.70	479.10
			105.6-167.1	168.1-246.9	260.8-879.9
3	0.38	0.92	81.96	171.9	756
			38.30-175.4	86.93-339.9	94.44-6052
5	0.23	0.94	31.25	85.1	631
			11.07-88.19	45.42-159.4	85.22-4672
7	0.24	0.93	8.21	23.45	191.12
			1.33-50.44	9.58-57.38	37.85-965.6

C. Effect on weight and volume of body

Measurements of the whole body weight of *C. longiareolata* larvae showed that the weight was affected under treatment of *L. nobilis* EO at its LC₂₅ and LC₅₀ (Table 3). The treatment decreased significantly the weight of larvae on the 3rd (p = 0.0041) and 7th day (p = 0.0096) after treatment with the highest dose.

However, treatment with *L. nobilis* essential oil caused a significant decrease in the body volume in larvae treated with the two concentrations (LC₂₅ and LC₅₀ respectively) at 3rd (p = 0.0023 and p = 0.0031) and 7th day (p = 0.0328 and 0.0094) after treatment. The body size is a pivotal trait for mosquitoes, because it can influence their blood-feeding ability, host attack rate and fecundity. All of these traits are important determinants of their potential to transmit diseases [46].

Dris et al. (2017) and Dris (2018) [25, 47] found that the morphometric parameters were decrease at different developmental stages of *C. pipiens* treated with *Ocimum basilicum* and *Mentha piperita* and *Lavandula dentata* respectively.

TABLE III. EFFECT OF *L. NOBILIS* EO (LC₂₅ AND LC₅₀) ON THE FRESH BODY WEIGHT (MG) AND BODY VOLUME (MM³) IN THE FOURTH INSTAR LARVAE OF *C. LONGIAREOLATA* (M ± SD, N=3 POOLS EACH CONTAINING 20 INDIVIDUALS)

Parameters	Time (Days)	Control	LC ₂₅	LC ₅₀
Volume (mm ³)	1	22.19 ± 4.45 A	16.48 ± 2.32 A	14.63 ± 1.91 A
	3	19.94 ± 5.44 A	14.14 ± 2.12 A	14.19 ± 2.45 B
	5	16.6 ± 2.21 A	14.14 ± 2.12 A	12.92 ± 2.16 A
	7	13.64 ± 2.13 A	12.83 ± 1.87 B	12.83 ± 1.87 C
Weight (mg)	1	6.055 ± 1.22 A	5.89 ± 0.76 B	4.79 ± 0.07 C
	3	5.96 ± 0.95 A	5.79 ± 0.27 B	4.69 ± 0.41 C
	5	5.94 ± 1.72 A	3.96 ± 1.30 B	2.91 ± 0.08 C
	7	5.04 ± 0.42 A	3.86 ± 0.06 A	2.82 ± 0.14 A

The different letters indicate significant differences among concentration treatments based on student test (p < 0.05).

D. Effect on biochemical composition

The amounts of carbohydrates, lipids and proteins were estimated in the whole body from different periods of fourth instar larvae of *C. pipiens* using LC₂₅ and LC₅₀ of *L. nobilis* essential oil (Table 4).

The comparison of mean values shows that a significant increase in the protein amounts was recorded in larvae at 1st (p = 0.0007 and p = 0.0002), 3rd (p = 0.0004 and p = 0.0003), 5th (p = 0.0002 and p < 0.001) and 7th day (p < 0.001). Concerning the carbohydrate levels, a significant reduction was observed at all tested periods after treatment with the highest dose (p = 0.0392, p = 0.0435, p = 0.0078 and p = 0.00215 respectively). Finally, the lipid content was decreased in treated series (LC₂₅ and LC₅₀) at 3rd day (p = 0.020 and 0.019), and only with the highest dose at 1st day (p = 0.0095) as compared to control series.

Several researches indicated that the exposure of an organism to xenobiotic product can modify the synthesis of certain proteins (enzymes of biotransformation, proteins of stress) [48]. Protein synthesis is necessary for the maintenance of body growth and reproduction [25]. Proteins enter in various reactions such as the hormonal regulation and they integrated in the cell as a structural element at the same time as the carbohydrates and the lipids [49; 50]. In the present investigation, after treatment of fourth instar larvae of *C. longiareolata* with *L. nobilis* EO, a stimulating action on proteins was generally exhibited during the tested periods. This stimulation might be due to the induction of protein synthesis such as some enzymatic and non-enzymatic biomarkers.

The reduction of the lipid levels after treatment with *L. nobilis* EO may be due to their effect on the lipid metabolism, and due to the utilization of lipid reserves for energy generation as a result of induced stress [51].

Carbohydrates play a crucial role in the physiology of the insects; the rates of glycogen in tissues are related to the physiological events such as the reproduction, the moult and the flight [52]. *Agastache foeniculum* EO applied in *Tribolium castaneum* reduced the energy reserves [53]. Askar et al. (2016) [54] reported that the application of clove oil in adults of three *Sitophilus* species increased lipid and protein levels, while

anise oil induce a reduction in the protein content in *S. granarius* and an increased in *S. oryzae*.

Many researchers have reported the depletion of larval energy reserves when exposed to several factors of stress (environmental, chemical and nutritional) [55- 56].

TABLE IV. EFFECT OF *L. NOBILIS* EO (LC25 AND LC50) ON AMOUNTS OF PROTEINS, CARBOHYDRATES AND LIPIDS (MG/INDIVIDUAL) FROM THE FOURTH INSTAR LARVAE OF *C. LONGIAREOLATA* AT DIFFERENT PERIODS (M ± SD, N = 3 POOLS EACH CONTAINING 20 INDIVIDUALS).

Content (µg/individual)	Time (Day)	Control	LC ₂₅	LC ₅₀
Proteines	1	139.11 ± 9.20 A	212.6 ± 0.75 B	239.6 ± 0.31 C
	3	133.77 ± 11.44 A	209.08 ± 3.67 B	217.1 ± 3.55 C
	5	128.8 ± 1.05 A	172.9 ± 3.80 B	220.8 ± 4.22 C
	7	116.16 ± 4.66 A	157.5 ± 2.17 B	219.68 ± 1.92 C
Lipids	1	68.05 ± 6.65 A	59.35 ± 9.15 A	43.21 ± 2.01 B
	3	67.35 ± 7.46 A	45.46 ± 2.04 B	43.14 ± 3.85 C
	5	52.82 ± 9.64 A	45.09 ± 2.14 A	40.26 ± 2.84 A
	7	44.36 ± 7.11 A	39.63 ± 0.76 A	38.42 ± 4.21 A
Carbohydrates	1	152.55 ± 12.08 A	126.56 ± 25.72 A	99.03 ± 19.54 B
	3	113.04 ± 13.71 A	107.01 ± 8.08 A	76.58 ± 6.81 B
	5	108.16 ± 7.19 A	93.43 ± 8.61 A	70.62 ± 7.13 B
	7	92.64 ± 5.92 A	82.80 ± 10.56 A	64.01 ± 8.02 B

IV. CONCLUSION

Phytoproducts possess different bioactive components that can be used as general toxicants against various larval stages of mosquitoes [57].

In the present study, it can be concluded that the essential oil of *L. nobilis* with Eucalyptol and Linalool as major compounds was found to exhibit potent larvicidal activity against *C. longiareolata* larvae. Moreover, the *L. nobilis* essential oil caused modification in morphometric parameters and biochemical composition.

ACKNOWLEDGMENT

This work was supported by the National Fund for Scientific Research to Pr. N. Soltani (Laboratory of Applied Animal Biology) and the Ministry of High Education and Scientific Research of Algeria (CNEPRU Project D01N01UN120120130005).

REFERENCES

[1] Chen S, Mulgrew B, Grant PM. A clustering technique for digital communications channel equalization using radial basis function networks. *IEEE Transactions on neural networks*. 1993; 4(4): 570-90. <https://doi.org/10.1109/72.238312>

[2] Edrissian GH. Malaria in Iran: Past and Present Situation. *Iranian Journal of Parasitology*. 2006; 1(1): 1-14.

[3] Khanavi M, Bagheri-Toulabi P, Abai MR, Sadati N, Hadjiakhoondi F, Hadjiakhoondi A, Vatandoost H. Larvicidal activity of marine algae,

Sargassum swartzii and *Chondria dasyphylla*, against malaria vector, *Anopheles stephensi*. *Journal of Vector Borne Diseases*. 2011; 48: 241–244.

[4] Francis S, Saavedra-Rodriguez K, Perera R, Paine M, Black IV WC, Delgoda R. Insecticide resistance to permethrin and malathion and associated mechanisms in *Aedes aegypti* mosquitoes from St. Andrew Jamaica. *PLoS one*. 2017 ; 26;12(6):e0179673.

[5] Mukandiwa L, Eloff JN and Naidoo V. Larvicidal activity of leaf extracts , seselin from *Clausena anisata* (Rutaceae) against *Aedes aegypti*. *South African Journal of Botany*. 2015 ; 1(100):169-73. <https://doi.org/10.1016/j.sajb.2015.05.016>

[6] Chandra G, Ghosh A, Bhattacharjee I, Ghosh SK. Use of larvivorous fish in biological and environmental control of disease vectors. *Biological and Environmental Control of Disease Vectors*. Wallingford (UK): CABI. 2013; 25-41.

[7] Ramírez-Lepe, Mario, Montserrat Ramírez-Suero. Biological control of mosquito larvae by *Bacillus thuringiensis* subsp. israelensis. *Insecticides-Pest Engineering*. InTech, 2012.

[8] Hamaidia K, Tine-Djebbar F, Soltani N. Activity of a selective insecticide (methoxyfenozide) against two mosquito species (*Culex pipiens* and *Culiseta longiareolata*): toxicological, biometrical and biochemical study. *Physiological Entomology*. 2018 ; DOI: 10.1111/phen.12261

[9] Tine-Djebbar F. & Soltani N. Activité biologique d'un agoniste non stéroïdien de l'hormone de mue sur *Culiseta longiareolata*: analyses morphométrique, biochimique et énergétique. *Synthèse*. 2008 ; 18 : 23–34.

[10] Rocha D, Novo M, Matos O, Figueiredo AC, Delgado M, Cabral MD, Liberato M, Moiteiro C. Potential of *Mentha pulegium* for mosquito control. *Revista de Ciências Agrárias*. 2015 ; 38 (2) : 155-65.

[11] El Akhal F, Greche H, Ouazzaoui CF, Guemmouh R, El Ouali Lalami A. Chemical composition and larvicidal activity of *Culex pipiens* essential oil of *Thymus vulgaris* grown in Morocco. *Journal of Materials and Environmental Science*. 2015; 1: 214-9.

[12] Benelli G, Pavela R, Canale A, Mehlhorn H. Tick repellents and acaricides of botanical origin: a green roadmap to control tick-borne diseases?. *Parasitology Research*. 2016; 115 (7): 2545–2560. <https://doi.org/10.1007/s00436-016-5095-1>

[13] Pavela R. Larvicidal effects of various Euro-Asiatic plants against *Culex quinquefasciatus* Say. larvae (Diptera: Culicidae). *Parasitology Research*. 2008; 102: 555–559. <https://doi.org/10.1007/s00436-007-0821-3>

[14] Geetha R and Roy A. Essential oil repellents: a short review. *International Journal of Drug Development and Research*. 2014; 6(2): 20-7

[15] Bakkali F, Averbeck S, Averbeck D, Idaomar M. Biological effects of essential oils—a review. *Food and chemical toxicology*. 2008; 1;46(2):446-75.

[16] Aurelie FD, Ascension NM, Gabriel TH, Herman AA, Pauline NM, Lebel TJ. Chemical composition and ovicidal, larvicidal and pupicidal activity of *Ocimum basilicum* essential oil against *Anopheles gambiae*. (Diptera: Culicidae). *European Journal of Medicinal Plants*. 2016; 16: 1-3. <https://doi.org/10.9734/EJMP/2016/28832>

[17] Aruna P, Murugan K, Priya A, Ramesh S. Larvicidal, pupicidal and repellent activities of *Gaultheria* oil (Plantae: Ericaceae) against the filarial vector, *Culex quinquefasciatus* (Insecta: Diptera: Culicidae). *Journal of Entomology and Zoology Studies*. 2014; 2: 290-4.

[18] Al-Sarar AS, Hussein HI, Abobakr Y. Chemical composition, adulticidal and repellent activity of essential oils from *Mentha longifolia* L. and *Lavandula dentata* L. against *Culex pipiens* L. *Mansoura Journal of Plant Protection and Pathology*, 2014; 5(7): 817-26.

[19] Aguiar RW, dos Santos SF, da Silva Morgado F, Ascencio SD, de Mendonca Lopes M, Viana KF, et al. Insecticidal and repellent activity of *Siparuna guianensis* Aubl. (Negramina) against *Aedes aegypti* and *Culex quinquefasciatus*. *PLoS One*. 2015; 10(2): 1-14. <https://doi.org/10.1371/journal.pone.0116765>

[20] Burt S. Essential oils: their antibacterial properties and potential applications in foods—a review. *International journal of food microbiology*. 2004 ; 94(3): 223-53. <https://doi.org/10.1016/j.ijfoodmicro.2004.03.022>

[21] Stefanello ME and Pascoal AC, Salvador MJ. Essential oils from neotropical Myrtaceae: chemical diversity and biological properties. *Chemistry & biodiversity*. 2011 ; 8(1):73-94 <https://doi.org/10.1002/cbdv.201000098>

- [22] Sukumar K, Perich MJ, Boobar LR. Botanical derivatives in mosquito control. A review. *Journal of the American Mosquito Control Association*. 1991 ; 7(2): 210-37.
- [23] Bouguerra N, Tine-Djebbar F, Soltani N. Algerian *Thymus vulgaris* essential oil: chemical composition and larvicidal activity against the mosquito *Culex pipiens*. *International Journal of Mosquito Research*. 2017; 4(1): 37-42.
- [24] Bouguerra N, Tine-Djebbar F, Soltani N. Effect of *Thymus vulgaris* L. (Lamiaceae) essential oil on energy reserves and biomarkers in *Culex pipiens* L. (Diptera: Culicidae) from Tebessa (Algeria). *Journal of Essential Oil Bearing Plants*. 2018; 21(4): 1082-95. <https://doi.org/10.1080/0972060X.2018.1504696>
- [25] Dris D, Tine-Djebbar F, Soltani N. Lavandula dentata essential oils: chemical composition and larvicidal activity against *Culiseta longiareolata* and *Culex pipiens* (Diptera: Culicidae). *African Entomology*. 2017; 25(2): 387-94. <https://doi.org/10.4001/003.025.0387>
- [26] El SN, Karagozlu N, Karakaya S, Sahn S. Antioxidant and antimicrobial activities of essential oils extracted from *Laurus nobilis* L. leaves by using solvent-free microwave and hydrodistillation. *Food and Nutrition Sciences*. 2014 ; 5 : 97-106. <https://doi.org/10.4236/fns.2014.52013>
- [27] Roselló J, Sempere F, Sanz-Berzosa I, Chiralt A, Santamarina MP. Antifungal activity and potential use of essential oils against *Fusarium culmorum* and *Fusarium verticillioides*. *Journal of Essential Oil Bearing Plants*. 2015 ; 18(2): 359-67. <https://doi.org/10.1080/0972060X.2015.1010601>
- [28] Ouibrahim A, Tlili-Ait-kaki Y, Bennadja S, Amrouni S, Djahoudi AG, Djebbar MR. Evaluation of antibacterial activity of *Laurus nobilis* L., *Rosmarinus officinalis* L. and *Ocimum basilicum* L. from Northeast of Algeria. *African journal of microbiology research*. 2013; 7(42): 4968-73. <https://doi.org/10.5897/AJMR2012.2390>
- [29] Akgül A, Kivanc M, Bayrak A. Chemical composition and antimicrobial effect of Turkish laurel leaf oil. *Journal of Essential Oil Research*. 1989 ; 1(6): 277-80.. <https://doi.org/10.1080/10412905.1989.9697798>
- [30] Tine-Djebbar F, Bouabida H, Soltani N. Répartition spatio-temporelle des Culicidés dans la région de Tébéssa. Editions Universitaires Européennes, 2016; 978(3) : 639-50856-7.
- [31] Jennings W, Shibamoto T. Qualitative analysis of flavour and fragrance volatile by glass capillary gas chromatography. New York Academic Press, 1980.
- [32] Adams RP. Identification of essential oil components by gas chromatograph/mass spectrometry. Carol Stream, USA: Allured Publishing Corporation, 1995.
- [33] Soltani-Mazouni N, Hami M, Gramdi H. Sublethal effects of methoxyfenozide on reproduction of the Mediterranean flour moth, *Ephestia kuehniella* Zeller. *Invertebrate reproduction & development*. 2012 ; 56(2): 157-63. <https://doi.org/10.1080/07924259.2011.582695>
- [34] Shibko S, Koivistoinen P, Tratnyek CA, Newhall AR, Friedman L. A method for sequential quantitative separation and determination of protein, RNA, DNA, lipid, and glycogen from a single rat liver homogenate or from a subcellular fraction. *Analytical biochemistry*. 1967; 19(3): 514-28. [https://doi.org/10.1016/0003-2697\(67\)90242-4](https://doi.org/10.1016/0003-2697(67)90242-4)
- [35] Bradford MM. A rapid, sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Analytical biochemistry*. 1976 ; 72(1-2): 248-54. [https://doi.org/10.1016/0003-2697\(76\)90527-3](https://doi.org/10.1016/0003-2697(76)90527-3)
- [36] Duchâteau G, Florquin M. Sur la tréhalosémie des insectes et sa signification. *Archives Internationales de Physiologie et de Biochimie*. 1959 ; 67(2): 306-14. <https://doi.org/10.3109/13813455909074435>
- [37] Goldsworthy GJ, Mordue W, Guthkelch J. Studies on insect adipokinetic hormones. *General and Comparative Endocrinology*. 1972 ; 18(3):545-51. [https://doi.org/10.1016/0016-6480\(72\)90034-2](https://doi.org/10.1016/0016-6480(72)90034-2)
- [38] Caputo L, Nazzaro F, Souza LF, Aliberti L, De Martino L, Fratianni F, Coppola R, De Feo V. *Laurus nobilis*: Composition of essential oil and its biological activities. *Molecules*. 2017 ; 22(6): 930. <https://doi.org/10.3390/molecules22060930>
- [39] Taoudiat A, Djenane D, Ferhat Z, Spigno G. The effect of *Laurus nobilis* L. essential oil, different packaging systems on the photo-oxidative stability of Chemlal extra-virgin olive oil. *Journal of Food Science and Technology*. 2018 ; 55(10): 4212-4222. <https://doi.org/10.1007/s13197-018-3357-x>
- [40] Da Silveira SM, Bittencourt LF, Fronza N, Cunha A, Neudt Scheuermann G, Werneck Vieira CR. Chemical composition and antibacterial activity of *Laurus nobilis* essential oil towards foodborne pathogens and its application in fresh Tuscan sausage stored at 7 °C. *LWT- Food Science Technology*. 2014 ; 59: 86–93. <https://doi.org/10.1016/j.lwt.2014.05.032>
- [41] Özcan, M., Chalchat, J. C. Effect of different locations on the chemical composition of essential oils of laurel (*Laurus nobilis* L.) leaves growing wild. Turkey. *Journal of Medicinal food*. 2005 ; 8(3) : 408-411. <https://doi.org/10.1089/jmf.2005.8.408>
- [42] Ben Jemâa MJ, Tersim N, Taleb Toudert K, Khouja ML. Insecticidal activities of essential oils from leaves of *Laurus nobilis* L. from Tunisia, Algeria and Morocco, and comparative chemical composition. *Journal of Stored Products Research*. 2012 ; 48 (97) :104.
- [43] Regnault-Roger C, Vincent C, Arnason JT. Essential oils in insect control: low-risk products in a high-stakes world. *Annual Review of Entomology*. 2012 ; 7(57): 405-24. <https://doi.org/10.1146/annurev-ento-120710-100554>
- [44] Dias CN, Moraes DF. Essential oils and their compounds as *Aedes aegypti* L. (Diptera: Culicidae) larvicides: review. *Parasitology Research*. 2014; 113: 565-92. <https://doi.org/10.1007/s00436-013-3687-6>
- [45] Venkatchalam MR, Jebanesan A. Larvicidal activity of *Hydrocotyle javanica* Thunb (Apiaceae) extract against *Culex quinquefasciatus*. *Journal of Experimental Zoology, India*. 2001; 4(1): 99–101.
- [46] Farjana T, Tuno N. Multiple blood feeding and host-seeking behavior in *Aedes aegypti* and *Aedes albopictus* (Diptera: Culicidae). *Journal of medical entomology*. 2013 ; 50(4): 838-46. <https://doi.org/10.1603/ME12146>
- [47] Dris D. larvicidal activity of the extracts of three plants: *Mentha piperita*, *Lavandula dentata* and *Ocimum basilicum* against larvae two species of mosquitoes *Culex pipiens* (Linné) and *Culiseta longiareolata* (Aitken). Doctoral Thesis. University of Badji Mokhtar, Annaba. 2018.
- [48] Curtis J, Omiecinski, John P, Vanden Heuvel, Gary H, Perdew, Jeffrey M. Peters. Xenobiotic metabolism, disposition, and regulation by receptors: from biochemical phenomenon to predictors of major toxicities. *Toxicological Sciences*. 2010 ; 120(1) : 49–75.
- [49] Cohen E. Chitin biochemistry: synthesis, hydrolysis and inhibition. *Advances in Insect Physiology*. 2010; 38: 5-74. [https://doi.org/10.1016/S0065-2806\(10\)38005-2](https://doi.org/10.1016/S0065-2806(10)38005-2)
- [50] Sugumaran M. Chemistry of cuticular sclerotization. *Advances in Insect Physiology*. 2010; 39: 151-209. <https://doi.org/10.1016/B978-0-12-381387-9.00005-1>
- [51] Sak O, Uckan F, Ergin E. Effects of cypermethrin on total body weight, glycogen, protein and lipid contents of *Pimpla turionellae* L. (Hymenoptera: Ichneumonidae). *Belgian Journal of Zoology*. 2006 ; 136: 53-58.
- [52] Kaufmann C.C, Brown C. Regulation of carbohydrate metabolism and flight performance by a hyper trehalosaemic hormone in the mosquito *Anopheles gambiae*. *Journal of Insect Physiology*. 2008 ; 54: 367-377. <https://doi.org/10.1016/j.jinsphys.2007.10.007>
- [53] Ebadollahi A, Khosravi R, Sendi JJ, Honarmand P and Amini RM. Toxicity and Physiological Effects of Essential Oil from *Agastache foeniculum* (Pursh) Kuntze Against *Tribolium castaneum* Herbst (Coleoptera: Tenebrionidae) Larvae. *Annual Review & Research in Biology*. 2013 ; 3(4): 649-658.
- [54] Askar SI, Al-Assaal MS, Nassar AMK. Efficiency of some essential oils and insecticides in the control of some *Sitophilus* insects (Coleoptera: Curculionidae). *Egyptian Journal of Plant Protection Research*. 2016 ; 4(2): 39-55.
- [55] Preet S, Sneha A. Biochemical evidence of efficacy of potash alum for the control of dengue vector *Aedes aegypti* (Linnaeus). *Parasitology Research*. 2011; 108: 1533-1539. <https://doi.org/10.1007/s00436-010-2210-6>
- [56] Vinayagam A, Senthilkumar N, Umamaheswari A. Larvicidal activity of some medicinal plant extracts against malaria vector *Anopheles stephensi*. *Research Journal of Parasitology*. 2008; 3: 50-58. <https://doi.org/10.3923/jp.2008.50.58>
- [57] Shaalan EA, Canyon D, Younes MW, Abdel-Wahab H, Mansour AH. A review of botanical phytochemicals with mosquitocidal potential. *Environment international* 2005 ; 31(8) : 1149-1166. <https://doi.org/10.1016/j.envint.2005.03.003>